


1. For the reaction $\text{BrO}_3^- + 5\text{Br}^- + 6\text{H}^+ \rightarrow 3\text{Br}_2 + 3\text{H}_2\text{O}$ at a particular time, $-\Delta[\text{BrO}_3^-]/\Delta t = 1.5 \times 10^{-2} \text{ M/s}$. What is $-\Delta[\text{Br}^-]/\Delta t$ at the same instant?

A) 13 M/s
 B) $7.5 \times 10^{-2} \text{ M/s}$
 C) $1.5 \times 10^{-2} \text{ M/s}$
 D) $3.0 \times 10^{-3} \text{ M/s}$
 E) 330 M/s

2. For the following reaction, $\Delta P(\text{C}_6\text{H}_{14})/\Delta t$ was found to be $-6.2 \times 10^{-3} \text{ atm/s}$.




Determine $\Delta P(\text{H}_2)/\Delta t$ for this reaction at the same time.

A) $6.2 \times 10^{-3} \text{ atm/s}$
 B) $1.6 \times 10^{-3} \text{ atm/s}$
 C) $2.5 \times 10^{-2} \text{ atm/s}$
 D) $-1.6 \times 10^{-3} \text{ atm/s}$
 E) $-2.5 \times 10^{-2} \text{ atm/s}$

3. The reaction $\text{A} + 2\text{B} \rightarrow \text{products}$ has the rate law, $\text{rate} = k[\text{A}][\text{B}]^3$. If the concentration of B is doubled while that of A is unchanged, by what factor will the rate of reaction increase?

A) 2 B) 4 C) 6 D) 8 E) 9


4. Consider the following reaction



If $[\text{C}]$ is increasing at the rate of $4.0 \text{ mol L}^{-1}\text{s}^{-1}$, at what rate is $[\text{B}]$ changing?

A) $-0.40 \text{ mol L}^{-1}\text{s}^{-1}$
 B) $-2.5 \text{ mol L}^{-1}\text{s}^{-1}$
 C) $-4.0 \text{ mol L}^{-1}\text{s}^{-1}$
 D) $-6.4 \text{ mol L}^{-1}\text{s}^{-1}$
 E) None of these choices is correct, since its rate of change must be positive.

5. For the reaction




the following data was collected at constant temperature. Determine the correct rate law for this reaction.

Trial	Initial [A] (mol/L)	Initial [B] (mol/L)	Initial Rate (mol/(L·min))
1	0.200	0.100	6.00×10^{-2}
2	0.100	0.100	1.50×10^{-2}
3	0.200	0.200	1.20×10^{-1}
4	0.300	0.200	2.70×10^{-1}

A) $\text{Rate} = k[\text{A}][\text{B}]$
 B) $\text{Rate} = k[\text{A}][\text{B}]^2$
 C) $\text{Rate} = k[\text{A}]^3[\text{B}]^2$
 D) $\text{Rate} = k[\text{A}]^{1.5}[\text{B}]$
 E) $\text{Rate} = k[\text{A}]^2[\text{B}]$

6. For the reaction



the following data was collected at constant temperature. Determine the correct rate law for this reaction.

Trial	Initial [A] (mol/L)	Initial [B] (mol/L)	Initial Rate (mol/(L·min))
1	0.125	0.200	7.25
2	0.375	0.200	21.75
3	0.250	0.400	14.50
4	0.375	0.400	21.75

A) $\text{Rate} = k[\text{A}][\text{B}]$
 B) $\text{Rate} = k[\text{A}]^2[\text{B}]$
 C) $\text{Rate} = k[\text{A}][\text{B}]^2$
 D) $\text{Rate} = k[\text{A}]$
 E) $\text{Rate} = k[\text{A}]^3$

7. The rate constant for a reaction is $4.65 \text{ L mol}^{-1} \text{ s}^{-1}$. What is the overall order of the reaction?
- Zero
 - First
 - Second
 - Third
 - More information is needed to determine the overall order.
8. Sulfuryl chloride, $\text{SO}_2\text{Cl}_2(g)$, decomposes at high temperature to form $\text{SO}_2(g)$ and $\text{Cl}_2(g)$. The rate constant at a certain temperature is $4.68 \times 10^{-5} \text{ s}^{-1}$. What is the order of the reaction?
- Zero
 - First
 - Second
 - Third
 - More information is needed to determine the overall order.
9. The data below were determined for the reaction $\text{S}_2\text{O}_8^{2-} + 3\text{I}^- (\text{aq}) \rightarrow 2\text{SO}_4^{2-} + \text{I}_3^-$.
- | Expt. # | $[\text{S}_2\text{O}_8^{2-}]$ | $[\text{I}^-]$ | Initial Rate |
|---------|-------------------------------|----------------|----------------------------------|
| 1 | 0.038 | 0.060 | $1.4 \times 10^{-5} \text{ M/s}$ |
| 2 | 0.076 | 0.060 | $2.8 \times 10^{-5} \text{ M/s}$ |
| 3 | 0.076 | 0.030 | $1.4 \times 10^{-5} \text{ M/s}$ |
- The rate law for this reaction must be:
- $\text{rate} = k[\text{S}_2\text{O}_8^{2-}][\text{I}^-]^3$
 - $\text{rate} = k[\text{S}_2\text{O}_8^{2-}]$
 - $\text{rate} = k[\text{S}_2\text{O}_8^{2-}]^2[\text{I}^-]^2$
 - $\text{rate} = k[\text{I}^-]$
 - $\text{rate} = k[\text{S}_2\text{O}_8^{2-}][\text{I}^-]$
10. At 25°C the rate constant for the first-order decomposition of a pesticide solution is $6.40 \times 10^{-3} \text{ min}^{-1}$. If the starting concentration of pesticide is 0.0314 M , what concentration will remain after 62.0 min at 25°C ?
- $1.14 \times 10^{-1} \text{ M}$
 - 47.4 M
 - $-8.72.0 \text{ M}$
 - $2.11 \times 10^{-2} \text{ M}$
 - $2.68 \times 10^{-2} \text{ M}$
11. A certain first-order reaction $\text{A} \rightarrow \text{B}$ is 25% complete in 42 min at 25°C . What is the half-life of the reaction?
- 21 min
 - 42 min
 - 84 min
 - 120 min
 - 101 min
12. A certain first-order reaction $\text{A} \rightarrow \text{B}$ is 25% complete in 42 min at 25°C . What is its rate constant?
- $6.8 \times 10^{-3} \text{ min}^{-1}$
 - $8.3 \times 10^{-3} \text{ min}^{-1}$
 - $3.3 \times 10^{-2} \text{ min}^{-1}$
 - $-3.3 \times 10^{-2} \text{ min}^{-1}$
 - 11 min^{-1}

Answers:

- Ans: B
- Ans: C
- Ans: D
- Ans: B
- Ans: E
- Ans: D
- Ans: C
- Ans: B
- Ans: E
- Ans: D
- Ans: E
- Ans: A